


Novel System for Immobilizing Foreign Proteins on the Cell Surface of *Bacillus subtilis*



BGSC Accession: ECE197 (see also ECE196 and 1A857)

Original Code: DH5 α (pNDH10)

Reference: Nguyen, H. D., and W. Schumann. 2006. Establishment of an experimental system allowing immobilization of proteins on the surface of *Bacillus subtilis* cells. *J. Biotechnol.* **122**:473-482.

Sequence: Not in database; available from BGSC at <http://www.bgsc.org/sequences/pNDH10.htm>

Features:

<i>xylR</i>	Repressor of the <i>B. megaterium xyl</i> operon; allows expression of <i>fbp</i> and fused sequences to be xylose-inducible
<i>fbpB</i>	C-terminal part of fibronectin binding protein B, including the sorting motif recognized by sortase A. Proteins fused to this fragment will be anchored to the cell wall of <i>B. subtilis</i> hosts that express SrtA.
<i>repA</i>	Initiation protein for theta-form Gram-positive plasmid replication; originally from pBS72.
<i>cat</i>	encodes chloramphenicol acetyl transferase; selectable in either <i>E. coli</i> or <i>B. subtilis</i> (chloramphenicol, 5 μ g/ml)
<i>bla</i>	encodes β -lactamase; selectable in <i>E. coli</i> only (ampicillin, 100 μ g/ml)

Description: pNDH10 is a shuttle vector, replicating at high copy number in *E. coli* in lower copy number in *Bacillus subtilis*. It features a xylose-inducible expression cassette upstream from the coding sequence of an anchoring protein that allows fused proteins to be anchored to the cell surface of a suitable host.

Construction: pNDH10 was constructed by inserting the 5' end of the fibronectin binding protein B gene (*fnbB*) of *Staphylococcus aureus* into the xylose-inducible expression vector pHCMC04.

Use: A coding sequence inserted inframe with the plasmid *fbp* gene fragment will have xylose-inducible expression. The fusion protein will be anchored on the cell surface of *B. subtilis* strain NDH03 (=BGSC 1A857) or any other host that expresses sortase A. Plasmid pNDH09 is an integrative vector that can be used to construct such a host in the desired genetic background.

Our thanks to Wolfgang Schumann for donating pNDH09, pNDH10, *B. subtilis* NDH03 and to the BGSC Collection!